

Zeng-Fu Xu · Whei-Lan Teng · Mee-Len Chye

## Inhibition of endogenous trypsin- and chymotrypsin-like activities in transgenic lettuce expressing heterogeneous proteinase inhibitor SaPIN2a

Received: 29 April 2003 / Accepted: 6 October 2003 / Published online: 23 October 2003  
© Springer-Verlag 2003

**Abstract** SaPIN2a, a proteinase inhibitor II from American black nightshade (*Solanum americanum* Mill.) is highly expressed in the phloem and could be involved in regulating proteolysis in the sieve elements. To further investigate the physiological role of SaPIN2a, we have produced transgenic lettuce (*Lactuca sativa* L.) expressing SaPIN2a from the *CaMV35S* promoter by *Agrobacterium*-mediated transformation. Stable integration of the *SaPIN2a* cDNA and its inheritance in transgenic lines were confirmed by Southern blot analysis and segregation analysis of the R<sub>1</sub> progeny. *SaPIN2a* mRNA was detected in both the R<sub>0</sub> and R<sub>1</sub> transformants on northern blot analysis but the SaPIN2a protein was not detected on western blot analysis using anti-peptide antibodies against SaPIN2a. Despite an absence of significant inhibitory activity against bovine trypsin and chymotrypsin in extracts of transgenic lettuce, the endogenous trypsin-like activity in each transgenic line was almost completely inhibited, and the endogenous chymotrypsin-like activity moderately inhibited. Our finding that heterogeneously expressed SaPIN2a in transgenic lettuce inhibits plant endogenous protease activity further indicates that SaPIN2a regulates proteolysis, and could be potentially exploited for the protection of foreign protein production in transgenic plants.

**Keywords** Chymotrypsin · *Lactuca* · Protease · Proteolysis · *Solanum* · Trypsin

**Abbreviations** *CaMV* cauliflower mosaic virus · *cDNA* complementary DNA · *NOS* nopaline synthase · *PAGE*

Z. Xu · W. Teng · M. Chye (✉)  
Department of Botany, The University of Hong Kong,  
Pokfulam Road, Hong Kong, China  
E-mail: mlchye@hkuce.hku.hk  
Fax: +852-28583477

*Present address:* Z. Xu  
Key Laboratory of Gene Engineering  
of the Ministry of Education, Zhongshan (Sun Yat-sen) University,  
Guangzhou 510275, China

polyacrylamide gel electrophoresis · *PI* proteinase inhibitor · *SaPIN2a* *Solanum americanum* proteinase inhibitor IIa · *SDS* sodium dodecyl sulphate · *T-DNA* transferred DNA

### Introduction

It has been proposed that plant proteinase inhibitors (PIs) which act on animal or microbial proteases play a role in the inhibition of proteolytic enzymes from pests or pathogens (Ryan 1981, 1989; Brzin and Kidric 1995). These conclusions arise from studies that use commercially available proteases, e.g. trypsin, chymotrypsin, elastase, and subtilisin from animal or microbial sources, as test enzymes in activity assays. However, none of these proteases is likely to be the true physiological target of these plant PIs (Laskowski and Kato 1980; Brzin and Kidric 1995).

Reports on the developmental regulation and tissue-specific accumulation of plant PIs (Rosahl et al. 1986; Sanchez-Serrano et al. 1986; Margossian et al. 1988; Hendriks et al. 1991; Pena-Cortes et al. 1991; Lorberth et al. 1992) do suggest they have endogenous functions. A soybean cysteine proteinase inhibitor has been designated a novel role in modulating programmed cell death (Solomon et al. 1999). Another, from cucumber leaves, which does not significantly inhibit commercial proteases of animal or microbial origin, inhibits cucumber glutamyl endopeptidase, suggesting regulation of its activity (Yamauchi et al. 2001).

Recently, we have shown that a *Solanum americanum* proteinase inhibitor II, SaPIN2a, is highly expressed in phloem (Xu et al. 2001). The localization of *SaPIN2a* mRNA and protein to the companion cells and sieve elements suggests a role in the regulation of proteolysis in phloem development/function. To further investigate, we have expressed SaPIN2a in transgenic lettuce and, here, we show that its heterogeneous expression results in the inhibition of plant endogenous protease activity.

Lettuce was chosen for expression of SaPIN2a because it neither possesses detectable trypsin inhibitory activity in its leaves nor responds to any treatments by accumulating inhibitors (Walker-Simmons and Ryan 1977).

## Materials and methods

### Plant material and growth conditions

Seeds of lettuce (*Lactuca sativa* L. cv. Great Lakes No.118) were obtained from Northrup King Co., Minneapolis, MN, USA. Plant tissue cultures were maintained and propagated in vitro in a growth chamber at 22–24 °C under a 12 h light/12 h dark regime. Plants in soil were grown under natural conditions in a greenhouse.

### Generation of transgenic lettuce plants expressing SaPIN2a

The *Agrobacterium tumefaciens*-mediated transformation vector pSa7 was constructed by replacing the  $\beta$ -glucuronidase (*GUS*) gene (Jefferson et al. 1987) in pBI121 (Clontech) with the *SaPIN2a* cDNA (Xu et al. 2001). To confirm the absence of any spurious ATG codon between the transcription start site (Jefferson et al. 1987) and the *SaPIN2a* start codon, this region was sequenced with a primer (5'-CAATCCCACTATCCTTCGCAAGACC-3') located within the cauliflower mosaic virus 35S (*CaMV35S*) promoter. The binary vector pSa7 was introduced into *A. tumefaciens* LBA4404 by direct transformation (Holsters et al. 1978). Lettuce was transformed according to Curtis et al. (1994) with modifications: petunia nurse cell cultures were omitted from the callus-inducing medium and kanamycin sulphate (100 µg/ml) was added to the media for callus induction, shoot regeneration and rooting.

### Segregation analysis of the progeny (R<sub>1</sub>) plants of transgenic lettuce

The segregation ratios of kanamycin-resistant (Km<sup>R</sup>) to kanamycin-sensitive (Km<sup>S</sup>) plants in the progeny (R<sub>1</sub>) of self-fertilized primary (R<sub>0</sub>) transgenic lettuce were determined by germinating surface-sterilized seeds from each R<sub>0</sub> line on Murashige and Skoog (MS) medium (Murashige and Skoog 1962) containing kanamycin sulphate (100 µg/ml). After incubation for 2–3 weeks in a tissue-culture chamber (22–24 °C, 12 h light/12 h dark), the seedlings were scored for kanamycin resistance. The segregation ratios were assessed by Chi-square analysis.

### Southern blot analysis

Twenty µg DNA, isolated (Dellaporta et al. 1983) from lettuce leaves, was digested with restriction endonucleases, separated by electrophoresis in a 0.8% agarose gel and blotted onto Hybond-N membrane (Amersham) according to Sambrook et al. (1989). The blot was pre-hybridized in 30% deionized formamide, 6× standard saline citrate (1× SSC: 150 mM NaCl, 15 mM sodium citrate, pH 7.0), 5× Denhardt's, 1% SDS, 50 µg/ml denatured, sonicated salmon sperm DNA at 42 °C for 4 h. Random-primed <sup>32</sup>P-labelled *SaPIN2a* cDNA was added for hybridization overnight at 42 °C. The blot was washed in 0.1× SSC, 0.1% SDS at room temperature.

### Northern blot analysis

Total RNA was extracted (Nagy et al. 1988) from nightshade (*Solanum americanum* Mill.) plants, wild-type lettuce or transgenic lettuce and analyzed by northern blot analysis as previously

described (Xu et al. 2001). The *SaPIN2a* cDNA was used in the generation of a random-primed <sup>32</sup>P-labelled probe.

### Western blot analysis

Total plant protein was extracted according to the procedure of Wu et al. (1997). Protein concentration was determined following Bradford (1976). Total protein was separated by 4–20% gradient SDS-PAGE (Gallagher 1995) for western blot analysis (Sambrook et al. 1989) using polyclonal antibodies raised in rabbit against a synthetic peptide (GESDPRNPKDC) corresponding to amino acids 77–87 of *SaPIN2a* (Xu et al. 2001). The Amplified Alkaline Phosphatase Immun-Blot Assay Kit (Bio-Rad) was used in detection of cross-reacting bands.

### Assay of trypsin and chymotrypsin inhibitory activities

Total plant proteins, extracted with 50 mM Tris (pH 8.1), 20 mM CaCl<sub>2</sub>, were used in spectrophotometric assays of trypsin or chymotrypsin inhibitory activity as described by Kollipara and Hymowitz (1992).

In the trypsin inhibitory-activity assay, 150 µl of leaf extract was pre-incubated for 3 min at room temperature (RT) in a quartz cuvette (10 mm path length, 3.5 ml capacity) with 100 µl of bovine trypsin (10 µg/ml in 1 mM HCl; Calbiochem Cat. No. 6502) and assay buffer [46 mM Tris-HCl (pH 8.1), 11.5 mM CaCl<sub>2</sub>], to give a final volume of 1.5 ml. The reaction was initiated by the addition of 1.5 ml of substrate [2 mM *p*-toluenesulphonyl-L-arginine methyl ester (TAME; Sigma Cat. No. T4626) in assay buffer] to the pre-incubation mixture. Recording of absorbance at 247 nm (A<sub>247</sub>) was immediately initiated. The spectrophotometer was set to auto-zero just before the start of recording and absorbance was measured at 30-s intervals for 3 min. In the standard reaction, 100 µl of bovine trypsin (10 µg/ml in 1 mM HCl), in the absence of leaf extract, was pre-incubated with 1.4 ml of assay buffer.

In the chymotrypsin inhibitory-activity assay, 50 µl of leaf extract was pre-incubated for 3 min at RT with 100 µl of bovine  $\alpha$ -chymotrypsin (20 µg/ml in 1 mM HCl; Calbiochem Cat. No. 230832) and assay buffer [0.1 M Tris-HCl (pH 7.8), 0.1 M CaCl<sub>2</sub>], to give a final volume of 1.5 ml. The reaction was initiated by the addition of 1.5 ml of substrate [1 mM N-benzoyl-L-tyrosine ethyl ester (BTEE; Sigma Cat. No. B6125) in 50% (w/w) methanol] to the pre-incubation mixture. Absorbance at 256 nm (A<sub>256</sub>) was similarly monitored as described above. In the standard reaction, 100 µl of chymotrypsin (20 µg/ml in 1 mM HCl), in the absence of leaf extract, was pre-incubated with 1.4 ml of assay buffer.

### Assay of endogenous trypsin- and chymotrypsin-like activities


Endogenous trypsin- and chymotrypsin-like activities in leaves of wild-type and transgenic R<sub>1</sub> lettuce plants were determined using the same procedures as in the trypsin and chymotrypsin inhibitory-activity assays, with the omission of bovine trypsin or chymotrypsin from the reaction.

## Results

### Southern blot analysis of transgenic lettuce plants

Lettuce was transformed using an *A. tumefaciens* LBA4404 derivative harboring plasmid pSa7 (Fig. 1a), the binary vector expressing the *SaPIN2a* cDNA from the *CaMV35S* promoter. Southern blot analyses were used to confirm the integration of the *SaPIN2a* cDNA

**Fig. 1a–c** The binary vector pSa7 used in lettuce (*Lactuca sativa*) transformation and Southern blot analysis of R<sub>0</sub> transgenic plants. **a** Structure of T-DNA region of pSa7. RB, right border of the transferred-DNA (T-DNA); NOS-Pro, nopaline synthase (NOS) promoter; NPT II, neomycin phosphotransferase II gene encoding resistance to kanamycin (Kan<sup>R</sup>); NOS-ter, NOS terminator; CaMV35S-Pro, promoter of cauliflower mosaic virus (CaMV) 35S RNA; SaPIN2a, *Solanum americanum* proteinase inhibitor IIa; LB, left border of the T-DNA. **b, c** Twenty µg of genomic DNA was digested with *Eco*RI (**b**) or *Bam*HI (**c**) and fractionated on a 0.8% agarose gel. The DNA was blotted and probed to <sup>32</sup>P-labelled random-primed



in the lettuce genome and to estimate the copy number of *SaPIN2a* cDNA in the various transgenic lines.

Results from Southern blot analysis using *Eco*RI-digested DNA of transgenic and regenerated (R<sub>0</sub>) lettuce lines, TL1, 7, 11, 15 and 33, show the presence of an expected 0.58-kb *Eco*RI band (indicated by an arrow in Fig. 1b), suggesting integration of the *SaPIN2a* cDNA in the lettuce genome. This band was lacking in wild-type lettuce (Fig. 1b, lane WT). Additional strong hybridization bands in TL7 and 33 may be due to incomplete *Eco*RI digestion of genomic DNA or the presence of rearranged copies of the *SaPIN2a* cDNA in these lines.

Since only one *Bam*HI site is present between the 35S promoter and the *SaPIN2a* cDNA within the transferred DNA (T-DNA) region on pSa7 (Fig. 1a), the number of bands hybridizing to <sup>32</sup>P-labelled *SaPIN2a* cDNA on Southern blot analysis of *Bam*HI-digested DNA should give a good estimate of the transgene copy number. Different hybridization patterns seen with *Bam*HI-digested DNA (Fig. 1c) implied that the various transgenic lines resulted from independent transformation events. Also, the number of strongly hybridizing bands ranged from one (TL1, 7, 11) to three (TL33), corresponding to single or multiple copies of transgene in these lines. Single copies of *SaPIN2a* cDNA in TL1 and 11, and multiple copies in TL33, were further confirmed by segregation analysis of their progenies (Table 1).

#### Expression of SaPIN2a mRNA in transgenic lettuce

The transcription of *SaPIN2a* in R<sub>0</sub> and R<sub>1</sub> transgenic lettuce lines was examined by northern blot analysis. Total RNA from leaves of R<sub>0</sub> transgenic plants, identified by Southern blot analysis (Fig. 1), was

hybridized to <sup>32</sup>P-labelled *SaPIN2a* cDNA. As shown in Fig. 2a, *SaPIN2a* mRNA was detected in the transgenic lines TL1, 7, 11, 15 and 33, but not in wild-type lettuce. The *SaPIN2a* transcript in transgenic lettuce (0.93 kb, indicated by an arrow in Fig. 2a) was slightly larger than the endogenous transcript of 0.67 kb in *S. americanum* (indicated by an arrowhead in Fig. 2a, lane Sa), due to the additional 0.26 kb from the nopaline synthase (NOS) terminator (Fig. 1a). We observed a more complex mRNA expression pattern in the R<sub>1</sub> transgenic lines, with the presence of a shorter transcript below the expected 0.93-kb band (Fig. 2b). It is worth noting that the shorter transcript found in leaves of some R<sub>0</sub> plants (Fig. 2a) became more prominent in leaves of R<sub>1</sub> plants (Fig. 2b).

Non-detection of SaPIN2a protein in transgenic lettuce by western blot analysis


Total leaf proteins from R<sub>0</sub> transgenic lettuce, expressing *SaPIN2a* mRNA on northern blot analysis (Fig. 2a), were

**Table 1** Segregation of the NPT-II gene (kanamycin resistance) in the progeny of transgenic lettuce (*Lactuca sativa*) plants

Parent transgenic line (R <sub>0</sub> )	Progeny (R <sub>1</sub> ) plants <sup>a</sup>		Expected ratio <sup>b</sup> (Kan <sup>R</sup> :Kan <sup>S</sup> )	Chi-square	Probability
	Kan <sup>R</sup>	Kan <sup>S</sup>			
TL1	67	21	3:1	0.061	1.000
TL11	30	10	3:1	0.000	1.000
TL33	176	3	255:1 63:1	7.587 0.015	0.667 1.000

<sup>a</sup>Kan<sup>R</sup> = kanamycin-resistant; Kan<sup>S</sup> = kanamycin-sensitive

<sup>b</sup>Ratio 3:1 is for a single functional locus, 255:1 for four functional loci and 63:1 for three functional loci




**Fig. 2a–d** Northern blot and western blot analyses on transgenic lettuce plants. For northern blot analysis (**a**, **b**), the blots were probed with random-primed  $^{32}\text{P}$ -labelled *SaPIN2a* cDNA. The hybridization band corresponding to the expected *SaPIN2a* transcript is indicated by an *arrowhead* (0.67-kb mRNA in *S. americanum* stems) or an *arrow* (0.93-kb mRNA in transgenic lettuce). The 25S ribosomal RNA (rRNA) bands stained with ethidium bromide are shown (*bottom panel*) to indicate the amount of total RNA loaded per lane. **a** Northern blot analysis of RNA from  $R_0$  transgenic lettuce lines. Total RNA (20  $\mu\text{g}$ ) was isolated from leaves of wild-type (*WT*) or transgenic lettuce and from *S. americanum* (*Sa*) stems as a positive control. **b–d** Total RNA (20  $\mu\text{g}$ ) and total protein (20  $\mu\text{g}$ ) were isolated from identical leaves and stems of 38-day  $R_1$  transgenic (*TL1*, *TL11*, *TL33*) and wild-type (*WT*) lettuce plants. Total RNA (20  $\mu\text{g}$ ) and total protein (14  $\mu\text{g}$ ) isolated from *S. americanum* stems (*Sa*) were included as positive controls. **b** Northern blot analysis of RNA from  $R_1$  transgenic lettuce plants. **c** Western blot analysis of protein from  $R_1$  transgenic lettuce plants using *SaPIN2a*-specific antibodies. Cross-reacting bands in lettuce leaves (18.1 kDa) and *S. americanum* stems (16.7 kDa) are indicated. **d** Coomassie blue stain of total protein from transgenic and wild-type plants separated on a 4–20% gradient SDS-PAGE gel to demonstrate amounts of total protein loaded in **c**

used in western blot analysis with affinity-purified *SaPIN2a*-specific antibodies. However, *SaPIN2a* could not be detected in the leaf proteins from these  $R_0$  transgenic lines (data not shown). Western blot analysis was also carried out using total proteins from  $R_1$  plants from three self-pollinated  $R_0$  lines (*TL1*, *11* and *33*). Figure 2c shows the results of western blot analysis on these  $R_1$  plants, using tissue samples identical to those used in the northern blot shown above (Fig. 2b). Despite very high levels of *SaPIN2a* mRNA in leaves of these transgenic lines (Fig. 2b), no protein band corresponding to *SaPIN2a* in *S. americanum* stem (Fig. 2c, lane *Sa*) was detected in transgenic lettuce leaf proteins (Fig. 2c). Some non-specific cross-reacting bands were observed and one (18.1 kDa), present in lettuce leaves but not stems, is very close to the size of native *SaPIN2a* (16.7 kDa; Fig. 2c). Since native *SaPIN2a* in *S. americanum* accumulates in stem (Xu et al. 2001), we were interested to investigate whether the cellular transport of *SaPIN2a* may have accounted for its apparent absence in the transgenic leaves. Hence, northern blot and western blot analyses were carried out using stems from these transgenic lettuce plants. Although *SaPIN2a* mRNA was expressed in their stems (Fig. 2b) at lower amounts than leaves, *SaPIN2a* protein remained undetected on western blots with *SaPIN2a*-specific antibodies (Fig. 2c).

#### Assay of trypsin and chymotrypsin inhibitory activities

Despite verification of the *SaPIN2a* sequence in pSa7 by DNA sequence analysis and confirmation of the integration of its cDNA by Southern blot analysis (Fig. 1b, c) and of its mRNA expression by northern blot analysis (Fig. 2a, b), the *SaPIN2a* protein was not observed on western blot analysis. The relatively low sensitivity of western blot analysis may have resulted in the non-detection of *SaPIN2a* in these transgenic lines. Hence, to further examine for the presence of *SaPIN2a*, we used proteinase inhibitory-activity assays, which we hoped would prove more sensitive than western blot analysis.

Crude leaf extracts from  $R_1$  transgenic and wild-type plants were tested for inhibitory activity against bovine trypsin and chymotrypsin. Results of trypsin inhibitory-activity assays (Fig. 3a) showed no significant inhibitory activity against bovine trypsin in transgenic lettuce. Leaf extracts from all three transgenic lines analyzed (*TL1*, *11*, *33*) showed trypsin activity similar to that of the standard reaction. Surprisingly, leaf extracts of wild-type plants had much higher trypsin activity than the standard reaction containing bovine trypsin (Fig. 3a). In the case of assay of chymotrypsin inhibitory activity (Fig. 3b), reactions of leaf extracts from both transgenic and wild-type plants showed higher chymotrypsin activity than the standard; each transgenic line showed slightly decreased activity compared to wild-type. These results of the inhibitory-activity assays suggest that these transgenic lettuce leaves may possess considerable endogenous trypsin- and chymotrypsin-like activities.



#### Assay of endogenous trypsin- and chymotrypsin-like activities

The endogenous trypsin- and chymotrypsin-like activities in leaves of wild-type and transgenic R<sub>1</sub> lettuce plants were determined using the same procedures as in the assays of proteinase inhibitory activity, with the omission of bovine trypsin or chymotrypsin from the reaction. Results of these experiments are shown in

**Fig. 3a–d** Assays of trypsin- and chymotrypsin inhibitory activities and trypsin- and chymotrypsin-like activities on transgenic lettuce leaf extracts. Leaf extracts were prepared from 54-day-old wild-type (*WT*) or transgenic (*TL1*, *TL11*, *TL33*) R<sub>1</sub> plants. Each value represents the mean  $\pm$  SE of three replicates. **a** Trypsin inhibitory-activity assay. One  $\mu$ g of bovine trypsin (Calbiochem) was incubated with 150  $\mu$ l of assay buffer (standard) or leaf extracts and the residual trypsin activity was determined by measuring the increase of absorbance at 247 nm during substrate hydrolysis. **b** Chymotrypsin inhibitory-activity assay. Two  $\mu$ g of bovine  $\alpha$ -chymotrypsin (Calbiochem) was incubated with 50  $\mu$ l of assay buffer (standard) or leaf extract and the residual chymotrypsin activity was determined by measuring the increase of absorbance at 256 nm during substrate hydrolysis. **c** Trypsin-like activity assay. Leaf extract (150  $\mu$ l) was directly incubated with substrate in the absence of bovine trypsin. Trypsin-like activity was determined by measuring the increase in absorbance at 247 nm during substrate hydrolysis. **d** Chymotrypsin-like activity assay. Leaf extract (50  $\mu$ l) was directly incubated with substrate in the absence of bovine  $\alpha$ -chymotrypsin. Chymotrypsin-like activity was determined by measuring the increase in absorbance at 256 nm during substrate hydrolysis

Fig. 3c, d. Both endogenous trypsin- and chymotrypsin-like activities were detected in leaf extracts from wild-type lettuce. However, the assay results from transgenic leaves were unexpected and interesting. The endogenous trypsin-like activities in all three transgenic plants were almost completely inhibited (Fig. 3c), while the endogenous chymotrypsin-like activities in these lines decreased moderately (Fig. 3d). The inhibition of endogenous trypsin- and chymotrypsin-like activities in transgenic lettuce leaves suggests that SaPIN2a protein accumulated at amounts that could inhibit endogenous trypsin- and chymotrypsin-like activities, despite being undetected on western blot analysis.

#### Discussion

We have demonstrated that the heterogeneous expression of a plant PI, SaPIN2a, in transgenic lettuce inhibits endogenous trypsin- and chymotrypsin-like activities. The significant inhibition of trypsin-like activity and the moderate inhibition of chymotrypsin-like activity in transgenic lettuce, resulting from the expression of SaPIN2a, could not be due to a mutation caused by the T-DNA insertion because all three independent lines (TL1, 11, 33) tested showed similar inhibition. This finding further supports our previous hypothesis that SaPIN2a has an endogenous role in regulating the activity of endogenous proteases in the phloem (Xu et al. 2001). Since it has been shown that the yield and quality of antibodies produced in transgenic plants are significantly affected by endogenous proteolytic degradation (Stevens et al. 2000), heterogeneous expression of PIs could be exploited in the protection of foreign protein production in transgenic plants. The identification of the target endogenous proteases for SaPIN2a would be our future goal. So far, only a trypsin-like enzyme and its endogenous inhibitor have been identified in lettuce seeds (Shain

and Mayer 1965, 1968). In barley, an endogenous protease, a carboxypeptidase, has been shown to hydrolyse a number of ester substrates of trypsin and chymotrypsin (Mikola and Pietila 1972).

The difference in the levels of *SaPIN2a* mRNA accumulation in leaves and stems of transgenic lettuce plants (Fig. 2b) was unexpected. The *CaMV35S* promoter is generally regarded as a strong constitutive promoter and directs high-level transcription in nearly all plant organs (Nagy et al. 1985; Odell et al. 1985). Our observations, taken together with those of others (Jefferson et al. 1987; Benfey et al. 1989; Williamson et al. 1989; Yang and Christou 1990; Sunilkumar et al. 2002), suggest that the *CaMV35S* promoter need not be constitutive.

It has been reported that the expression of transgenes in lettuce is not as efficient as in other transgenic plants (McCabe et al. 1999; Ryder 1999). The mechanism of low transgene expression in lettuce is not well understood and may be associated with DNA methylation (McCabe et al. 1999). Observations on the lack of foreign protein accumulation despite over-expression of its corresponding mRNA have also been reported in transgenic petunia (Jones et al. 1985), tobacco (Jones et al. 1985; Florack et al. 1994), tomato (Seymour et al. 1993), cauliflower (Passelegue and Kerlan 1996) and potato (Gatehouse et al. 1997). In this study, we detected shorter transcripts other than that expected of the *SaPIN2a* mRNA using the *NOS* terminator in transgenic lettuce. This suggests that the mRNA may not have undergone proper processing and could have been further degraded. Inadvertently, this may be a contributing factor to the failure in the detection of SaPIN2a protein in transgenic lettuce on western blot analysis, despite high accumulation of the *SaPIN2a* mRNA.

Whether SaPIN2a possesses the putative inhibitory activity towards bovine trypsin and chymotrypsin, based on its sequence homology to known PIN2 proteins, is still unclear. One possibility is that the quantity of SaPIN2a protein in leaves of transgenic lettuce is insufficient, as indicated by western blot analysis, for assay of *in vitro* inhibitory activity using bovine trypsin or chymotrypsin. Nonetheless, this amount of SaPIN2a is sufficient to inhibit endogenous trypsin-like activity in lettuce leaves. An alternative explanation is that SaPIN2a could be specific to certain plant endogenous proteases but not to bovine trypsin and chymotrypsin. Protease inhibitors from soybean (Birk et al. 1963) and wheat (Applebaum and Konijn 1966) that inhibited larval gut proteolysis of *Tribolium castaneum* were inactive towards mammalian trypsin and chymotrypsin. The maize proteinase inhibitor (MPI), belonging to the potato proteinase inhibitor I (PIN1) family (Cordero et al. 1994), effectively inhibited midgut chymotrypsin of *Spodoptera littoralis* larvae, but only weakly inhibited bovine chymotrypsin, unlike most PIN1 members which are potent inhibitors of mammalian chymotrypsin (Tamayo et al. 2000). To unequivocally establish the

inhibitory activities of SaPIN2a, purified SaPIN2a protein from *S. americanum* stems should be employed for further *in vitro* activity assays.

**Acknowledgments** This work was supported by funds from The University of Hong Kong (to M.-L.C.). Z.-F.X. received a postgraduate studentship from The University of Hong Kong.

## References

- Applebaum SW, Konijn AM (1966) The presence of a *Tribolium*-protease inhibitor in wheat. *J Insect Physiol* 12:665–669
- Benfey PN, Ren L, Chua NH (1989) The CaMV 35S enhancer contains at least two domains which can confer different developmental and tissue-specific expression patterns. *EMBO J* 8:2195–2202
- Birk Y, Gertler A, Khalef S (1963) Separation of a *Tribolium*-protease inhibitor from soybeans on a calcium phosphate column. *Biochim Biophys Acta* 67:326–328
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein–dye binding. *Anal Biochem* 72:248–254
- Brzin J, Kidric M (1995) Proteinases and their inhibitors in plants: role in normal growth and in response to various stress conditions. *Biotechnol Genet Eng Rev* 13:420–467
- Cordero MJ, Raventos D, Segundo BS (1994) Expression of a maize proteinase inhibitor gene is induced in response to wounding and fungal infection: systemic wound-response of a monocot gene. *Plant J* 6:141–150
- Curtis IS, Power JB, Blackhall NW, de Laat AMM, Davey MR (1994) Genotype-independent transformation of lettuce using *Agrobacterium tumefaciens*. *J Exp Bot* 45:1441–1449
- Dellaporta SL, Wood J, Hicks JB (1983) A plant DNA mini-preparation: version II. *Plant Mol Biol Rep* 1:19–21
- Florack DEA, Dirkse WG, Visser B, Heidekamp F, Stiekema WJ (1994) Expression of biologically active hordeothionin in tobacco. Effects of pre- and pro-sequences at the amino and carboxyl termini of the hordeothionin precursor on mature protein expression and sorting. *Plant Mol Biol* 24:83–96
- Gallagher SR (1995) Separation of proteins on gradient gels. In: Coligan JE, Dunn BM, Ploegh HL, Speicher DW, Wingfield PT (eds) *Current protocols in protein science*, vol 1. Wiley, New York, pp 10.1.17–10.1.23
- Gatehouse AMR, Davison GM, Newell CA, Merryweather A, Hamilton WDO, Burgess EPJ, Gilbert RJC, Gatehouse JA (1997) Transgenic potato plants with enhanced resistance to the tomato moth, *Lacanobia oleracea*: growth room trials. *Mol Breed* 3:49–63
- Hendriks T, Vreugdenhil D, Stiekema WJ (1991) Patatin and four serine proteinase inhibitor genes are differentially expressed during potato tuber development. *Plant Mol Biol* 17:385–394
- Holsters M, de Waele D, Depicker A, Messens E, van Montagu M, Schell J (1978) Transfection and transformation of *Agrobacterium tumefaciens*. *Mol Gen Genet* 163:181–187
- Jefferson RA, Kavanagh TA, Bevan MW (1987) GUS fusions:  $\beta$ -glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J* 6:3901–3907
- Jones JDG, Dunsmuir P, Bedbrook J (1985) High level expression of introduced chimaeric genes in regenerated transformed plants. *EMBO J* 4:2411–2418
- Kollipara KP, Hymowitz T (1992) Characterization of trypsin and chymotrypsin inhibitors in the wild perennial *Glycine* species. *J Agric Food Chem* 40:2356–2363
- Laskowski M Jr, Kato I (1980) Protein inhibitors of proteinases. *Annu Rev Biochem* 49:593–626
- Lorberth R, Dammann C, Ebnet M, Amati S, Sanchez-Serrano JJ (1992) Promoter elements involved in environmental and developmental control of potato proteinase inhibitor II expression. *Plant J* 2:477–486

- Margossian LJ, Federman AD, Giovannoni JJ, Fischer RL (1988) Ethylene-regulated expression of a tomato fruit ripening gene encoding a proteinase inhibitor I with a glutamic residue at the reactive site. *Proc Natl Acad Sci USA* 85:8012–8016
- McCabe MS, Mohapatra UB, Debnath SC, Power JB, Davey MR (1999) Integration, expression and inheritance of two linked T-DNA marker genes in transgenic lettuce. *Mol Breed* 5:329–344
- Mikola J, Pietila K (1972) Hydrolysis of ester substrates of trypsin and chymotrypsin by barley carboxypeptidase. *Phytochemistry* 11:2977–2980
- Murashige T, Skoog F (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plant* 15:473–497
- Nagy F, Odell JT, Morelli G, Chua NH (1985) Properties of expression of the 35S promoter from CaMV in transgenic tobacco plants. In: Zaitlin M, Day P, Hollaender A (eds) *Biotechnology in plant science: relevance to agriculture in the eighties*. Academic Press, New York, pp 227–235
- Nagy F, Kay SA, Chua NH (1988) Analysis of gene expression in transgenic plants. In: Gelvin SB, Schilperoort RA (eds) *Plant molecular biology manual*. Kluwer, Dordrecht, pp B4:1–29
- Odell JT, Nagy F, Chua NH (1985) Identification of DNA sequences required for activity of the cauliflower mosaic virus 35S promoter. *Nature* 313:810–812
- Passelegue E, Kerlan C (1996) Transformation of cauliflower (*Brassica oleracea* var. *botrytis*) by transfer of cauliflower mosaic virus genes through combined cocultivation with virulent and avirulent strains of *Agrobacterium*. *Plant Sci* 113:79–89
- Pena-Cortes H, Willmitzer L, Sanchez-Serrano JJ (1991) Abscisic acid mediates wound induction but not developmental-specific expression of the proteinase inhibitor II gene family. *Plant Cell* 3:963–972
- Rosahl S, Eckes P, Schell J, Willmitzer L (1986) Organ-specific gene expression in potato: isolation and characterization of tuber-specific cDNA sequences. *Mol Gen Genet* 202:368–373
- Ryan CA (1981) Proteinase inhibitors. In: Marcus A (ed) *The biochemistry of plants*, vol 6. Academic Press, New York, pp 351–370
- Ryan CA (1989) Proteinase inhibitor gene families: strategies for transformation to improve plant defenses against herbivores. *BioEssays* 10:20–24
- Ryder EJ (1999) *Lettuce, endive and chicory*. CABI Publishing, New York
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
- Sanchez-Serrano JJ, Schmidt R, Schell J, Willmitzer L (1986) Nucleotide sequence of proteinase inhibitor II encoding cDNA of potato (*Solanum tuberosum*) and its mode of expression. *Mol Gen Genet* 203:15–20
- Seymour GB, Fray RG, Hill P, Tucker GA (1993) Down-regulation of two non-homologous endogenous tomato genes with a single chimaeric sense gene construct. *Plant Mol Biol* 23:1–9
- Shain Y, Mayer AM (1965) Proteolytic enzymes and endogenous trypsin inhibitor in germinating lettuce seeds. *Physiol Plant* 18:853–859
- Shain Y, Mayer AM (1968) Activation of enzymes during germination — trypsin-like enzyme in lettuce. *Phytochemistry* 7:1491–1498
- Solomon M, Belenghi B, Delledonne M, Menachem E, Levine A (1999) The involvement of cysteine proteases and protease inhibitor genes in the regulation of programmed cell death in plants. *Plant Cell* 11:431–443
- Stevens LH, Stoopen GM, Elbers IJW, Molthoff JW, Bakker HAC, Lommen A, Bosch D, Jordi W (2000) Effect of climate conditions and plant developmental stage on the stability of antibodies expressed in transgenic tobacco. *Plant Physiol* 124:173–182
- Sunilkumar G, Mohr L, Lopata-Finch E, Emani C, Rathore KS (2002) Developmental and tissue-specific expression of CaMV 35S promoter in cotton as revealed by GFP. *Plant Mol Biol* 50:463–474
- Tamayo MC, Rufat M, Bravo JM, Segundo BS (2000) Accumulation of a maize proteinase inhibitor in response to wounding and insect feeding, and characterization of its activity toward digestive proteinases of *Spodoptera littoralis* larvae. *Planta* 211:62–71
- Walker-Simmons M, Ryan CA (1977) Wound-induced accumulation of trypsin inhibitor activities in plant leaves. *Plant Physiol* 59:437–439
- Williamson JD, Hirsch-Wyncott ME, Larkins BA, Gelvin SB (1989) Differential accumulation of a transcript driven by the CaMV 35S promoter in transgenic tobacco. *Plant Physiol* 90:1570–1576
- Wu Y, Llewellyn D, Mathews A, Dennis ES (1997) Adaptation of *Helicoverpa armigera* (Lepidoptera: Noctuidae) to a proteinase inhibitor expressed in transgenic tobacco. *Mol Breed* 3:371–380
- Xu ZF, Qi WQ, Ouyang XZ, Yeung E, Chye ML (2001) A proteinase inhibitor II of *Solanum americanum* is expressed in phloem. *Plant Mol Biol* 47:727–738
- Yamauchi Y, Ejiri Y, Sugimoto T, Sueyoshi K, Oji Y, Tanaka K (2001) A high molecular weight glutamyl endopeptidase and its endogenous inhibitors from cucumber leaves. *J Biochem* 130:257–261
- Yang NS, Christou P (1990) Cell type specific expression of a CaMV 35S-GUS gene in transgenic soybean plants. *Dev Genet* 11:289–293